



# BMJ Open Influence of physical fitness components on personality factors and risk perception of children and adolescents: a cross-sectional study

Noelia González-Gálvez,<sup>1</sup> Raquel Vaquero-Cristobal ,<sup>2</sup> María José Maciá-Andreu,<sup>1</sup> Marta García-Tascon,<sup>3</sup> Antonio Soler-Marín ,<sup>4</sup> Ana María Gallardo-Guerrero<sup>1</sup>

**To cite:** González-Gálvez N, Vaquero-Cristobal R, Maciá-Andreu MJ, *et al*. Influence of physical fitness components on personality factors and risk perception of children and adolescents: a cross-sectional study. *BMJ Open* 2023;**13**:e071995. doi:10.1136/bmjopen-2023-071995

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2023-071995>).

Received 20 January 2023  
Accepted 27 October 2023



© Author(s) (or their employer(s)) 2023. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

<sup>1</sup>Facultad de Deporte, UCAM Universidad Católica de Murcia, Murcia, Región de Murcia, Spain

<sup>2</sup>Department of Physical Activity and Sport, Faculty of Sport Sciences, University of Murcia, Murcia, Región de Murcia, Spain

<sup>3</sup>Departamento del Deporte e Informática, Pablo de Olavide University, Sevilla, Andalucía, Spain

<sup>4</sup>Dpto. Tecnología de la Alimentación y Nutrición, UCAM Universidad Católica de Murcia, Murcia, Región de Murcia, Spain

## Correspondence to

Raquel Vaquero-Cristobal;  
[raquel.vaquero@um.es](mailto:raquel.vaquero@um.es)

## ABSTRACT

**Objectives** To examine the associations of cardiorespiratory fitness (VO<sub>2</sub> max) and muscular strength with indicators related to the risk scale, such as perceived competence, sensation seeking, competitiveness, risk taking and risk perception in sports.

**Design** Cross-sectional study.

**Setting** High schools from the Region of Murcia (Spain).

**Participants** Three-hundred-and-seventeen adolescents participated (mean age: 13.69±1.2 years old).

**Primary and secondary outcome measures** Body mass, body height, Course-Navette test, upper limb strength and psychoeducational factors that determine the propensity towards sports accidents in school children, the Sports Accident Propensity Scale were evaluated. It was performance t-test for independent samples, stepwise multiple linear regression models and a multiple mediation analysis.

**Results** The analysis showed significant differences with respect to sex in height, VO<sub>2</sub> max, handgrip strength and in all factors of the questionnaire (p=0.02-<0.01). Adolescents who presented greater VO<sub>2</sub> max, strength in the handgrip test and age showed a higher score in factors 1 and 3. Higher scores in factor 2 were associated with better VO<sub>2</sub> max and strength in handgrip test. Youngers and better values of strength in the handgrip showed higher score in factors 4 and 5. The mediation analysis with two mediating variables (handgrip strength and VO<sub>2</sub> max) showed a significant indirect effect. When handgrip strength and VO<sub>2</sub> max were included in the equations, the association between sex and each factor ceased to be significant.

**Conclusion** This study highlights the potential benefits of muscular strength (handgrip) and VO<sub>2</sub> max in the perceived risk scale, and the variable of age on this.

**Trial registration number** Clinical trial: NCT05544370 (pre-results).

## INTRODUCTION

Numerous benefits of physical fitness for physical, cognitive and psychological health are well known in children and adolescents<sup>1-3</sup> although in some cases, they may also entail a risk of suffering from some kind

## STRENGTHS AND LIMITATIONS OF THIS STUDY

- ⇒ The main strength of the present investigation was the possibility of carrying out a study relating the propensity of sports accidents with the physical condition of adolescents.
- ⇒ Face-to-face surveys were used, which made it possible to avoid the bias that commonly involves the use of technology, and all the doubts of the respondents were resolved.
- ⇒ It should be noted that the sample size was a limitation, as it would be interesting to increase sample size to obtain a higher representation of this sector.

of injury.<sup>4 5</sup> Roldán-Vendrell<sup>6</sup> indicates that risk is the possibility that a person will suffer a certain injury or accident.<sup>4 7</sup> Sport accidents can have a negative effect not only on athletes, but may also affect three areas, such as people, facilities and equipment or areas related to compliance with the regulations of the sport.<sup>8 9</sup>

A multidimensional analysis is required that takes into account physical, psychological and situational aspects, for the correct study of the causes responsible for sports injuries and accidents,<sup>10</sup> and up-to-date accident data records. Some studies in Spain indicate that injuries and accidents represent 0.8% of physical education (PE) classes, with a slightly higher incidence in males.<sup>11</sup> In Ireland, the study by O'Toole *et al*<sup>12</sup> conducted in two Irish paediatric orthopaedic outpatient departments, indicated that upper limb injuries were the most common type of injury in sports and recreation. Afterwards, the authors O'Connor *et al*<sup>13</sup> found that more than a quarter of school children were at risk of injury each year, and of those who sustained an injury, more than a third would sustain two or more injuries during the year, with the

most common injuries being strains, bruises, sprains and fractures.

Other aspects related to risk propensity are perceived competence, sensation seeking, competitiveness, risk taking and risk perception.<sup>14</sup> In general, Latorre states that children tend to underestimate risks, believe they are invulnerable and in some situations take significant risks in physical-sports practice,<sup>15</sup> making it necessary to strengthen measures to ensure not only the processes of purchasing equipment, but also to ensure their corresponding inspections, proper use and maintenance.

Regarding perceived competence, an athlete's low level of perceived competence was associated with a higher risk of injury,<sup>16</sup> while at the same time, it made them feel less competent for adopting risky behaviours,<sup>17</sup> with it being lower in females.<sup>18–24</sup> Likewise, a relationship has been observed between muscular strength and cardiovascular capacity, and fitness and perceived competence.<sup>25</sup> Regarding sensation seeking, athletes with lower levels of this factor are less likely to take risks,<sup>26</sup> and are also influenced by variables such as personality type, type of sport and gender.<sup>27–30</sup> On the other hand, being older is directly related to lower levels of sensation seeking, and it is therefore believed that this relationship may be related to biological processes.<sup>31</sup> In relation to competitiveness, it may be related to the behaviours that may be dangerous to physical integrity.<sup>10</sup> In addition, high levels of competitive anxiety are associated with an increased risk of injury,<sup>32</sup> and the studies confirmed that boys show higher values of this factor, although it was not related to age.<sup>33–35</sup> Horvath and Zuckerman<sup>36</sup> claim that successful past experiences with risk-taking can lead to a reduction in perceived risk. Finally, regarding the risk perception factor, Kontos<sup>16</sup> indicates that it is associated with a higher risk of injury depending on age, previous experience, perceived competence, personality and gender, with men perceiving less risk than women.<sup>16 17 37–40</sup> In contrast, no differences in risk perception in sports were found between boys and girls.<sup>41</sup>

Some studies have also analysed the relationship between these factors. Kontos<sup>17</sup> found that an overestimation of competence was positively related to risk-taking. On the other hand, sensation seeking influences risk perception, and in this case, risky behaviour and accident rates, being a clear predictor of risky behaviour.<sup>42</sup> With regard to the variables of age, gender and muscle strength in the hands, relationships between them were found in a study of 144 climbers from 22 different countries. The authors stated that grip strength in men is superior to that of women, but no significant differences were detected according to the age of the participants analysed.<sup>43</sup>

Without a doubt, special care must be taken when sports are practiced in facilities with children. As Latorre states, they do not yet have a sufficient capacity to discern and anticipate risks and/or dangers.<sup>15</sup> Furthermore, certain behavioural factors can contribute to unintentional injuries,<sup>44</sup> and even more so in school settings, as research discrepancies have been observed between maintaining

the balance between ensuring children's safety and providing them with physically and emotionally stimulating environments.<sup>45</sup> There are numerous studies on physical condition, maximum O<sub>2</sub> volume, muscle strength, body composition, among others in children and adolescents, but there is no research where these physical abilities are related to psychological aspects such as Sports Accident Propensity Scale, where variables such as risk perception are analysed. This approach is carried out in order to determine the profiles of the students, to raise awareness of the risks involved in physical and sporting activities at school<sup>40</sup> and to develop appropriate injury prevention strategies.<sup>46 47</sup> The aim of this study was to examine the associations between sex, age and physical fitness components, especially cardiorespiratory fitness and muscular strength, and indicators relative to the risk scale such as perceived competence, search for sensations, competitiveness, assumption of risk and risk perception in sports of youth/adolescents, in order to prevent sports-related accidents.

## MATERIAL AND METHODS

### Study design

This study was conducted in the Region de Murcia (Spain) (online supplemental file 1). Parents or guardians and adolescents were informed about the project, and an informed consent was obtained from all study participants and their parents/legal tutors (online supplemental file 2). All the data collected were treated confidentially and anonymously. The present investigation was conducted in accordance with the Declaration of Helsinki. This cross-sectional study design followed the Strobe Statement (online supplemental file 3). The data from this research are part of a registered clinical trial (NCT05544370 / pre-results).

### Sampling method and sample size

The participants were volunteer adolescents from two high schools from the Region de Murcia. They were aged between 12 and 17 years old (mean age: 13.69±1.2). The inclusion criteria for participating in the study were as follows: (a) being enrolled in one academic year in Secondary School, (b) being present on assessment day and (c) being authorised to participate in the measurements. The exclusion criteria were as follows: having any musculoskeletal, neurological, cardiovascular, metabolic or rheumatic alterations.

In order to establish the sample size and power, Rstudio V.3.15.0 software was used. The significance level was set to  $\alpha=0.05$ . According to the SD established for VO<sub>2</sub> max (mL/kg/min) in previous studies<sup>48</sup> and an estimated error of 1.5 mL/kg/min, a valid sample size of 313 was needed for a CI of 95%. Finally, the sample consisted of 317 adolescents, resulting in an error of 1.437 mL/kg/min. Males represented 48.37% of the sample, while the remaining 51.63% was female.

In addition, the body mass index (BMI) of the students in the sample was similar to those found in the results from the Health Behaviour in School-aged Children (HBSC) survey in Europe and Canada. This implies that the adolescents included in this study were representative of those included for adolescents in Spain as a whole.

### Procedures

The same trained researchers measured the variables using standardised conditions in a single session, from 9:00 am to 12:00 pm from May 2022 to June 2022. The adolescents were instructed to wear lightweight clothes. No warm-ups were performed before the evaluations. Before the examinations, to establish the reliability of the examiner, a double-blind study was performed with 30 participants, obtaining an intraclass correlation coefficient higher than 95%.

### Anthropometry measures

Body mass was measured using a SECA 762 scale (SECA, Germany) and height using a GPM anthropometer (Siber-Hegner, Switzerland). After this, BMI was calculated with the Quetelet Index formula ( $\text{BMI} = \text{body mass (kg)} / \text{height (m)}^2$ ).<sup>49</sup>

### Physical fitness measures

The Course-Navette test (a 20m Shuttle run test) was used to assess cardiorespiratory fitness (CRF) as described previously.<sup>50</sup> The maximum oxygen consumption ( $\text{VO}_2$  max, mL/kg/min) was estimated from the number of laps performed by the test participants using the equation reported by Leger *et al*<sup>50</sup> ( $\text{VO}_2 \text{ max} = 31.025 + 3.238 X - 3.248 A + 0.1536 A X$ , where  $X = \text{running speed}$  and  $A = \text{age}$ ).

To measure upper limb strength, the participants performed a handgrip strength test. Handgrip strength was measured two times on each arm in a standing position with the arms at the sides. Each participant was asked to squeeze the grip with maximal strength or 3s alternatively with right and left hands, with the elbow in full extension with 1 min rest between the measurements to avoid local muscle fatigue. The grip span of the dynamometer was adjusted according to the hand size of the adolescent. The maximum score in kilograms for each hand was recorded.<sup>51</sup> A digital grip strength dynamometer was used for this (TKK 5401; Takei Scientific Instruments, Tokyo, Japan).

### Psychological aspects measures

In order to assess the psychoeducational factors that determine the propensity of sports accidents in school children, the Sports Accident Propensity Scale (*Escala de Propensión al Accidente Deporte*, EPAD) was used. This scale is composed of 27 items that describe variables that condition the possibility of having a sports accident. This questionnaire presents a Likert-type scale where 1 means 'strongly disagree' and 6 means 'strongly agree'. It includes items from five different factors: perceived competence; sensation seeking; competitiveness; risk taking and risk perception.<sup>14</sup>

### Patient and public involvement

Patients and/or the public were not involved in the design, performance, reporting or dissemination plans of this research.

### Statistical analysis

The database for this research is available from the corresponding author and are publicly available.<sup>52</sup> Normality analyses were performed using the Kolmogorov-Smirnov test and Mauchly's W-test for sphericity analysis. To establish differences between sexes, a t-test for independent samples was used. The effect size was calculated with Cohen's coefficients. A value higher than 0.8 was considered a high effect, a value between 0.6 and 0.4 was considered a moderate effect, a value between 0.4 and 0.2 was considered as a low-moderate effect and a value lower than 0.2 was considered a low effect size.<sup>53</sup> The following was used to determine the variables to be included in the stepwise multiple linear regression models. To analyse whether there was a correlation between a categorical variable and a continuous variable, Spearman's correlation statistic was used. Pearson's correlation analysis was used to analyse the correlation between continuous variables. To analyse the association between categorical variables in the 2x2 tables, Cramer's V statistical test was applied: and in the 2xn tables, the contingency coefficient, showing the value of the statistic and the p-value. The maximum expected value was 0.707, with  $r < 0.3$  indicating a low association, a moderate association was defined as an  $r$  value between 0.3 and 0.5, and a high association as  $r > 0.5$ . Stepwise multiple linear regression models were used to predict each factor according to the variables. In the case that a non-linear multiple regression model provided the best explanation of the variance, as compared with the multiple linear regression models, the best model association between the dependent and independent variables was explored with a curvilinear estimation. The multiple mediation analysis, SPSS macro PROCESS (model 4), was applied with two independent significant mediators (SPSS, Chicago, IL, USA). A classical Baron and Kenny step regression method was used.<sup>54</sup> If the association between dependent and independent variables disappeared after the mediation variable was included, the mediation variable was considered as a complete mediator. The statistical analyses were performed with the IBM SPSS package (V.25.0). The significance level set at  $p < 0.05$ . In a complementary manner, a generalisability analysis was carried out to verify that the estimated results were reliable and generalisable, with the SAGT V.1.0 software.<sup>55</sup>

### RESULTS

The baseline characteristics of the adolescents are shown in [table 1](#). [Table 2](#) shows the data according to sex and the analysis of differences.

[Table 3](#) shows the linear regression of sex with the factors from the questionnaire. It shows that sex significantly

**Table 1** Basic characteristics of the adolescents (n=317)

Variables	Mean±SD/% (n)
Age (years old)	13.69±1.20
Sex	
Male	48.58 (154)
Female	51.42 (163)
Height (cm)	161.38±8.91
Body mass (kg)	54.09±11.32
BMI (kg/m <sup>2</sup> )	20.63±3.71
VO <sub>2</sub> max (mL/kg/min)	20.86±4.80
Handgrip (kg)	23.90±6.81
EPAD scale	
Factor 1. Perceived competence	28.73±9.37
Factor 2. Search for sensations	27.04±8.49
Factor 3. Competitiveness	13.71±5.21
Factor 4. Assumption of risk	11.54±5.21
Factor 5. Risk perception	10.86±4.75

%, percentage; BMI, body mass index; cm, centimetres; kg, kilograms; mL, millilitre; N, sample; Vo<sub>2</sub> VO<sub>2</sub> max, maximum oxygen uptake.

explained all the factors, although it only explained a small percentage of the variance (from 1 to 5.7%).

When incorporating the rest of the variables into the model (table 4), the stepwise multiple linear regression showed that the variables VO<sub>2</sub> max, handgrip strength and age, explained most of the factor values. Adolescents who presented a greater VO<sub>2</sub> max, greater strength in the handgrip test and greater age showed a higher score in factor 1 (perceived competence)

**Table 3** Linear regression of each factor of EPAD scale by sex

Variables	R <sup>2</sup>	P value	Standardised coefficients (β)
Factor 1. Perceived competence	0.06	<0.01	-0.24
Factor 2. Search for sensations	0.01	0.04	-0.20
Factor 3. Competitiveness	0.04	<0.01	-0.20
Factor 4. Assumption of risk	0.02	0.02	-0.14
Factor 5. Risk perception	0.03	0.01	-0.15

and factor 3 (competitiveness). In relation to factor 2 (sensation seeking), higher scores were associated with better VO<sub>2</sub> max and a greater strength in the handgrip test. The students who were younger and had a higher level of strength in the handgrip test showed a higher assumption of risk and a lower risk perception (factors 4 and 5).

The stepwise multiple linear regression analysis did not include sex. When the association between sex and the variables included in the model (handgrip strength and VO<sub>2</sub> max) was verified by means of a correlation, we proceeded to the mediation study. The mediation analysis with two mediation variables (handgrip strength and VO<sub>2</sub> max) showed a significant and indirect effect. When handgrip strength and VO<sub>2</sub> max were included in the equations, the association between sex and each factor was no longer significant (figure 1).

Finally, the analysis of generalisability (online supplemental files 4,5) shows in the first design, a generalisability coefficient of 0.902. This result shows the high reliability

**Table 2** Analysis of gender differences

Variables	Male (n=154) Mean±SD	Female (n=163) Mean±SD	Mean Diff±SD Diff Mean±SD	95% CI	F	P value	Effect size
Age (years old)	13.73±1.21	13.66±1.19	0.07±0.13	-0.19 to 0.32	1.28	0.61	0.06
Height (cm)	162.61±9.86	160.23±7.72	2.38±0.98	0.44 to 4.33	7.63	0.02	0.27
Body mass (kg)	54.41±10.61	53.91±12.03	0.70±0.50	1.27 to -2.01	3.00	0.70	0.04
BMI (kg/m <sup>2</sup> )	20.40±3.51	20.88±3.92	-0.48±0.04	-1.30 to 0.34	2.19	0.25	0.13
VO <sub>2</sub> max (mL/kg/min)	21.76±4.44	20.03±4.98	1.73±0.55	0.64 to 2.82	0.32	<0.01	0.37
Handgrip (kg)	25.66±7.70	22.12±5.31	3.54±0.74	2.09 to 5.00	19.66	<0.01	0.54
EPAD scale							
Factor 1. Perceived competence	30.99±9.03	26.53±9.22	4.46±1.03	2.43 to 6.50	0.02	<0.01	0.49
Factor 2. Search for sensations	28.08±8.67	26.06±8.24	2.02±0.96	0.13 to 3.91	0.46	0.04	0.24
Factor 3. Competitiveness	14.70±5.03	12.76±5.23	1.95±0.58	0.80 to 3.09	0.09	<0.01	0.38
Factor 4. Assumption of risk	12.25±5.53	10.84±4.81	1.41±0.59	0.26 to 2.57	3.05	0.02	0.27
Factor 5. Risk perception	11.59±4.71	10.14±4.70	1.02±0.42	0.19 to 1.85	0.07	0.02	0.31

%, percentage; BMI, body mass index; cm, centimetres; Diff, differences; kg, kilograms; mL, millilitre; N, sample; Vo<sub>2</sub> max, maximum oxygen uptake.

**Table 4** Stepwise multiple linear regression analysis of the relationship of each factor of the EPAD scale with independent variables

Dependent variable		R <sup>2</sup>	P value	Included independent variables	β	P value	
Factor 1. Perceived competence	Model 1	0.18	<0.01	VO <sub>2</sub> max	0.42	<0.01	
	Model 2	0.34	<0.01	VO <sub>2</sub> max	0.82	<0.01	
				Age	0.57	<0.01	
	Model 3	0.37	<0.01	VO <sub>2</sub> max	0.71	<0.01	
				Age	0.42	<0.01	
	Factor 2. Search for sensations				Handgrip	0.19	<0.01
Model 1		0.06	<0.01	VO <sub>2</sub> max	0.24	<0.01	
Model 2		0.11	<0.01	VO <sub>2</sub> max	0.24	<0.01	
				Handgrip	0.22	<0.01	
Factor 3. Competitiveness		Model 1	0.06	<0.01	Handgrip	0.25	<0.01
		Model 2	0.11	<0.01	Handgrip	0.24	<0.01
	VO <sub>2</sub> max				0.21	<0.01	
	Model 3	0.15	<0.01	Handgrip	0.16	0.13	
				VO <sub>2</sub> max	0.45	<0.01	
				Age	0.34	<0.01	
Factor 4. Assumption of risk	Model 1	0.08	<0.01	Handgrip	0.29	<0.01	
	Model 2	0.10	<0.01	Handgrip	0.34	<0.01	
				Age	-0.14	0.03	
Factor 5. Risk perception	Model 1	0.05	<0.01	Handgrip	0.22	<0.01	
	Model 2	0.06	<0.01	Handgrip	0.27	<0.01	
				Age	-0.13	0.04	

B, standardised coefficients.

of the test. In addition, the percentage of variance (see online supplemental file 5) was found to be high.


## DISCUSSION

The aim of this study was to examine the associations of sex, age and physical fitness components, especially CRF and muscular strength, with indicators relative to the risk scale, such as perceived competence, search for sensations, competitiveness, assumption of risk and risk perception in sports of youth/adolescents, in order to prevent sports accident.

With regard to the influence of sex on the indicators related to risk propensity, men had a statistically significant higher propensity in all the dimensions of the EPAD scale analysed as compared with the women, as in a previous study,<sup>14</sup> in which this variable was highlighted as one of the determinants of risk behaviour.<sup>56</sup> Specifically, in relation to perceived competence, adolescents with a high estimation of their abilities would be more likely to adopt risky behaviours, as compared with others with a low estimation,<sup>17</sup> with this being higher in men in almost all related studies,<sup>21 24 57–61</sup> although Kontos<sup>16</sup> found no sex differences in self-perceived ability. In this regard, men tend to overestimate their competence while the

opposite is true for women,<sup>62</sup> a fact that is also reflected during childhood, where overestimation of one's physical abilities represents a risk factor for everyday injuries, with boys being more at risk than girls.<sup>63</sup> Regarding sensation seeking, the present research coincided with other studies in highlighting higher values in men,<sup>14 24 64–67</sup> which was also accentuated with age.<sup>65 68 69</sup>

These differences may be driven by both biological and socialisation factors.<sup>31</sup> Those with lower levels of sensation seeking try to avoid taking risks,<sup>26</sup> so in this case, women may be less predisposed. In this regard, previous research highlights that boys more often attribute their injuries to bad luck, leading them to experience repetitive injuries, while girls attribute injuries to their own behaviours, causing them to alter their behaviour in future situations to avoid them.<sup>56 70</sup> In relation to competitiveness and the influence of gender, the level was higher in men than in women, with results similar to previous studies.<sup>10 33–35 67</sup> The results referring to risk-taking showed higher values in males,<sup>16 17 37</sup> and these differences may increase when considering physical skills.<sup>37</sup> This could be due to gender-specific sport socialisation, with females focusing their training on skill development, and males on strength development, physical preparation and aggressiveness.<sup>71 72</sup>



**Figure 1** Simple mediator effect of multiple mediators of two mediation variables.

During adolescence, individuals are more susceptible to risk-taking, and therefore, to injury, with the percentage being higher in males.<sup>17</sup> Moreover, at this stage, other individual characteristics such as age, experience or temperament<sup>56 73</sup> may influence children's decision-making and therefore participation in risky activities. Finally, regarding risk perception, it was higher in females than in males, as in previous research,<sup>16 17</sup> although some studies found no gender differences.<sup>41 74</sup> The differences found may be based on psychosocial factors associated with the predominant male stereotype for boys in sports, as girls are taught to be wary of risks and boys are taught to accept them.<sup>75</sup> This is reflected in a greater male liking for risky sports as compared with females.<sup>74</sup> However, risk perception is also affected by other internal factors such as age, previous experiences, perceived competence, and personality.<sup>40</sup>

Although most previous research focused on sex as a predictor of risk propensity, in the present study, a multiple regression analysis highlighted a significant indirect effect of other mediating variables (VO<sub>2</sub> max, handgrip strength and age), which explained most of the factors analysed. However, this may be due to biological sex differences during adolescence, which may influence young people's perceptions and behaviours,<sup>72</sup> with significantly higher VO<sub>2</sub> max<sup>76</sup> and handgrip strength values detected in males. More specifically, higher values of VO<sub>2</sub> max, handgrip strength and an older age, were related to higher perceived competence and greater competitiveness. In relation to sensation seeking, higher values were associated with better results in the VO<sub>2</sub> max value and handgrip strength test. Finally, younger adolescents with a higher strength in the handgrip test showed higher risk-taking and lower risk perception values. VO<sub>2</sub> max levels may therefore positively influence optimism, which plays an important role in self-regulation and adaptive behaviour in children and adolescents,<sup>77</sup> with higher VO<sub>2</sub> max values being associated with a greater perceived competence.<sup>58 78</sup>

Regarding handgrip strength, as a variable related to muscular strength, the results followed the line of previous benchmark research, in which muscular strength was associated to a higher self-esteem,<sup>2 58 79</sup> self-perception,<sup>2</sup> optimism<sup>79</sup> and perceived competence.<sup>58</sup> This is especially relevant during the adolescence, with substantial differences in the development and attainment of strength, agility and motor skills concomitant with variations in body size.<sup>80</sup>

Finally, age emerged as a predictor of perceived competence, competitiveness, risk-taking and risk perception. This evidence may be based on a possible direct relationship between biological factors, such as age and body size and risk behaviours in young athletes and adolescents,<sup>17</sup> with certain sociodemographic variables such as age being among determinants of risk behaviour.<sup>56 81</sup> and the detection of an increase in risk behaviours at ages 12–17 years old,<sup>17</sup> detecting within this stage that 15-year-olds show better inhibitory

control than 12-year-olds, while 17-year-olds show a better perspective than younger adolescents.<sup>82</sup> In this regard, López-Araujo and Osca<sup>83</sup> indicated that young people were characterised by underestimating the possibility of suffering certain risks, did not perceive them holistically, detected them more slowly, and tended to overestimate their skills.

Regarding risk-taking, Kern *et al*<sup>84</sup> found that among skateboarders, age was one of the significant explanatory variables, with young people being more likely to take risks than adults, as there was a sense of invincibility and greater impulsivity in youth than in adults. This could be conditioned by hormonal influences, in particular the increase in testosterone and oestradiol during adolescence, which are associated with an impulsive personality, increased risk-taking behaviour and impulsive personality.<sup>85</sup> The results obtained in several investigations are in line with the highlighted age-related differences in risk perception in sports, which makes adolescents more likely to take risks than adults.<sup>78–80</sup> At these ages, participants in school sport activities, depending on their perception of risk, will decide whether or not to take the risk, which is a relevant factor in explaining how they cope in these situations and a topic to be considered in the safety of the PE lessons.<sup>15</sup> Therefore, education for safety in sports should aim at ensuring that children know how to identify sources of risk, that they acquire an adjusted perception of the level of risk associated with it, and that they develop strategies to avoid them.<sup>86</sup> Furthermore, some studies highlight the idea that as children become older, their perception of their own motor competence becomes more accurate, and they therefore tend to have a more realistic estimate of their abilities.<sup>87</sup> In addition, fear caused by natural inhibition dependent on age and maturational stage is reduced as the child experiences a motivating emotion and learns to master appropriate challenges.<sup>88</sup> However, Kontos<sup>17</sup> and De Meester *et al*<sup>23</sup> indicated no significant differences in the perception of risk of injury in sports according to age. Similarly, the period from late childhood to adolescence is characterised by multiple physical, psychological and social changes,<sup>89</sup> with age influencing physical fitness, and with all the fitness components expected to improve from childhood to adolescence,<sup>90 91</sup> in particular, VO<sub>2</sub> max.<sup>92 93</sup>

With regard to the conclusions reached in accordance with the objectives of the present research, men have a higher propensity towards risk in all the variables analysed. However, sex was not found to be an explanatory factor for risk propensity when including VO<sub>2</sub> max, handgrip strength and age, which explains most of the factors analysed. In particular, a higher VO<sub>2</sub> max, a higher handgrip strength and being older showed higher scores in factor 1 (perceived competence) and factor 3 (competitiveness). In relation to factor 2 (sensation seeking), higher scores were associated with better VO<sub>2</sub> max and higher handgrip strength. Younger students with a higher level of strength in the handgrip test showed a higher assumption and a lower risk perception (factors 4 and 5).

However, there are other sources of variation, in addition to those studied in this research, which are closely related to the aforementioned constructs, and which should also be taken into account when preventing sports accidents. One of them is the type of sport practised, finding that those who choose to practice risky sports have higher levels of competitiveness, lower risk perception and greater sensation seeking.<sup>94 95</sup> Another factor to take into account is the personality of the subject. In this respect, previous research has associated a personality with low conscientiousness, low agreeableness, less cooperative and high neuroticism with higher risk-taking and participation in risky sports.<sup>26 96–100</sup> In addition, emotion dysregulation was also associated with increased risk-taking.<sup>101 102</sup> Finally, situational factors such as the influence of the group or the time urgency of the situation may distort the perception of the objectivity of the risk,<sup>15 103</sup> as well as the subject's previous successful experiences, reducing perceived risk and increasing risk-taking.<sup>36</sup>

This study is not without limitations. The cross-sectional design of this study is one of the first limitations that does not allow establishing causal relationships; thus, longitudinal studies are necessary. In addition, the method used to assess CRF was the Course-Navette test. Although this method is used in most studies, and is valid and reliable for children and adolescents,<sup>50</sup> this is an indirect method with limitations in its interpretation. The importance of the results obtained in the present research lies in the fact that a positive relationship between risk behaviours and injuries was found,<sup>17</sup> so it is necessary to investigate the variables that can influence them in order to prevent injuries during adolescence. In this regard, health-related fitness was the strongest predictor of future physical activity in the transition from primary to secondary school,<sup>104</sup> and given its relevance, its analysis as a possible predictor variable of risk behaviours is warranted.

## CONCLUSIONS

This study mainly highlights the potential benefits of muscular strength (handgrip) in all indicators of the risk perception scale, the CRF (VO<sub>2</sub> max) in three out of five components of the risk scale (ie, perceived competence, search for sensations and competitiveness), and the variable of age in all the factors, except for sensation seeking.

Therefore, it is essential to carry out actions aimed at the entire education community to raise awareness about the possible contingencies caused by poor planning, as well as training and awareness of good practices regarding the use of sports materials and equipment.<sup>7 82</sup>

Some strategies for the future could be as follows:

- ▶ To plan this work as a longitudinal study.
- ▶ To use go to another assess CRF assessment test to compare with the Course-Navette.
- ▶ To compare the results obtained which other risk perception tests.

- ▶ It is important to contact health centres to analyse the type of paediatric injuries and compare them with results from other countries.
- ▶ To conduct a pilot study on the analysis of children's injuries during one school year in different primary schools.

**Acknowledgements** The authors would like to thank the indispensable collaboration of the students and teachers of Physical Education from the Secondary School IES Dr Pedro Guillen and IES José Luis Castillo Puche. Thanks also to all the volunteers who helped with the data collection.

**Contributors** NG-G conceptualised. NG-G, MG-T and AMG-G designed the study. NG-G carried out the statistical analysis. NG-G and RV-C recruited the participants. NG-G, RV-C, MJM-A, AS-M and AMG-G collected the data. NG-G, AMG-G and RV-C organised the database. NG-G, RV-C, MJM-A, MG-T, AS-M and AMG-G wrote the first manuscript draft, the final manuscript draft, conducted the English proofreading and reviewed and edited the final version of the manuscript. NG-G is responsible for the overall content as guarantor. All authors contributed to the manuscript revision and approved the final version.

**Funding** This research was supported by two grant Research Support Plans from the Universidad Católica de Murcia (Spain), named 'H2PE: Promoting cardiometabolic health by 2 HIIT protocol, twice a week at Physical Education' (code: PMAFI-11/19); and named 'New technologies as tools for the promotion of health in schoolchildren of Compulsory Secondary Education of the CARM: effects of sports technology applications on the physical and psychological health of adolescents' (code: PMAFI-11/21).

**Competing interests** None declared.

**Patient and public involvement** Patients and/or the public were involved in the design, or conduct, or reporting or dissemination plans of this research. Refer to the Methods section for further details.

**Patient consent for publication** Consent obtained from parent(s)/guardian(s).

**Ethics approval** This study involves human participants and was approved by Ethics Committee of the Catholic University of Murcia (CE061914). Participants gave informed consent to participate in the study before taking part.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Data availability statement** Data are available in a public, open access repository. Technical appendix, statistical code and data set available from the Dryad repository, DOI: <https://doi.org/10.5061/dryad.v9s4mw70v>.

**Supplemental material** This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

## ORCID iDs

Raquel Vaquero-Cristobal <http://orcid.org/0000-0003-2708-4817>

Antonio Soler-Marín <http://orcid.org/0000-0002-1843-4005>

## REFERENCES

- 1 Ortega FB, Silventoinen K, Tynelius P, *et al*. Muscular strength in male adolescents and premature death: cohort study of one million participants. *BMJ* 2012;345:e7279.
- 2 Smith JJ, Eather N, Morgan PJ, *et al*. The health benefits of muscular fitness for children and adolescents: A systematic review and meta-analysis. *Sports Med* 2014;44:1209–23.

- 3 Prochnow T, Delgado H, Patterson MS, et al. Social network analysis in child and adolescent physical activity research: A systematic literature review. *J Phys Act Health* 2020;17:250–60.
- 4 Magaz-González AM, García-Tascón M. La Seguridad Deportiva a debate II. Madrid: Dykinson. In: Magaz-González AM, Flores-Allende G, García-Tascón M, eds. *Factores que afectan y agentes implicados en la seguridad deportiva*. 2022: 87–10.
- 5 Díaz-Pereira M, Buceta J, María A. Situaciones Estresantes Y Vulnerabilidad a Las Lesiones Deportivas: UN Estudio con Deportistas de Equipo. *Revista de Psicología Del Deporte* 2009;13.
- 6 Roldán Vendrell C. *Manual de Seguridad en los Centros Educativos*. Andalucía: Consejería de Educación y Ciencia. Junta de Andalucía, 2002.
- 7 Maciá MJ, Gallardo AM, Sánchez J, et al. Análisis de la Seguridad del Equipamiento Deportivo en Educación Secundaria Obligatoria. *Apunts* 2020:67–57.
- 8 Magaz-González AM. La Seguridad Deportiva a debate. Madrid: Dykinson. In: Magaz-González AM, Alias A, Jaenes JC, eds. *Perspectivas de tratamiento de la seguridad deportiva*. 2020: 13–30.
- 9 Meyerber M, Fraisse B, Dhalluin T, et al. Trampoline injuries compared with other child activities. *Arch Pediatr* 2019;26:282–4.
- 10 Latorre Román PÁ, Pantoja Vallejo A. Diseño Y Validación de UN Cuestionario de Propensión al Accidente Deportivo: (PAD-22). *CPD* 2013;13:51–62.
- 11 Ministerio de Sanidad y Consumo. Instituto Nacional de Consumo. In: *Programa de prevención de lesiones: Red de detección de accidentes domésticos y de ocio*. 2008.
- 12 O'Toole RV, Andersen RC, Vesnovsky O, et al. Are locking screws advantageous with plate fixation of Humeral shaft fractures? A Biomechanical analysis of synthetic and Cadaveric bone. *J Orthop Trauma* 2008;22:709–15.
- 13 O'Connor S, Whyte E, Chéilleachair NN. n.d. Sport and recreation musculoskeletal injuries in Irish primary school children.
- 14 Latorre-Román PÁ, Cámara-Pérez JC, Pantoja-Vallejo A, et al. Factores Psicoeducativos que Determinan La Propensión al Accidente Deportivo en Escolares: Diseño Y Validación de UN Cuestionario de Evaluación. *Analesps* 2013;29.
- 15 Latorre PA et al. Aspectos Comportamentales Y de Personalidad que Pueden Determinar La Propensión al Accidente Deportivo en Escolares. In: García-Tascón M, Magaz-González AM, Alias A, eds. *La Seguridad Deportiva a Debate*. Madrid: Dykinson, 2020: 153–68. Available: <https://www.dykinson.com/libros/la-seguridad-deportiva-a-debate/9788413246895/> [accessed 14 Dec 2022].
- 16 Kontos AP. Perceived risk, risk taking, estimation of ability and injury among adolescent sport participants. *J Pediatr Psychol* 2004;29:447–55.
- 17 Kontos AP. The effects of perceived risk of injury, risk-taking Behaviour, and body size on injury in youth sport. 2000.
- 18 Gentile A, Boca S, Giammusso I. You play like a woman! effects of gender stereotype threat on women's performance in physical and sport activities: A meta-analysis. *Psychology of Sport and Exercise* 2018;39:95–103.
- 19 Influence of gender on the tactical skill and motivational aspects in invasion sports in physical education. *CCD* 2019;14:93–105.
- 20 Corr M, McSharry J, Murtagh EM. Adolescent girls' perceptions of physical activity: A systematic review of qualitative studies. *Am J Health Promot* 2019;33:806–19.
- 21 Grandmontagne A, Ruiz de Azúa S, Liberal I. Propiedades Psicométricas de UN Nuevo Cuestionario para La Medida del Autoconcepto Físico. *Revista de Psicología Del Deporte* 2004;13.
- 22 Luis-de Cos G, Arribas-Galarraga S, Luis-de Cos I, et al. Motor competence, commitment, and anxiety in girls during physical education classes. *Retos* 2019:231–8.
- 23 De Meester A, Barnett LM, Brian A, et al. The relationship between actual and perceived motor competence in children, adolescents and young adults: A systematic review and meta-analysis. *Sports Med* 2020;50:2001–49.
- 24 Babí Lladós J, Inglés Yuba E, Cumellas Ruiz L, et al. El Perfil de Los Corredores Y su Propensión al Accidente Deportivo. *Revista Internacional de Medicina y Ciencias de La Actividad Física y Del Deporte* 2018;18:723.
- 25 Lloyd RS, Faigenbaum AD, Stone MH, et al. Position statement on youth resistance training: the 2014 International consensus. *Br J Sports Med* 2014;48:498–505.
- 26 Klina P, Burnik S, Kajtna T. Personality and sensation seeking in high-risk sports. *Acta Gymnica* 2017;47:41–8.
- 27 Zuckerman M. *Sensation seeking and risky behavior*. Washington: American Psychological Association, 2007.
- 28 Machado TS, Serrano JM, Silveira PL, et al. La Búsqueda de Sensaciones en Paracaidistas Portugueses. *Retos* 2022;44:1073–81.
- 29 Guskowska M, Boldak A. Sensation seeking in males involved in recreational high risk sports. *Biol Sport* 2010;27:157–62.
- 30 Burnik S, Jug S, Kajtna T. Sensation seeking in Slovenian female and male mountain climbers. *Acta Universitatis Palackianae Olomucensis* 2009;38.
- 31 Librán E. Búsqueda de Sensaciones. *Psicothema, ISSN 0214-9915* 2000;12:229–35.
- 32 Lavallée L, Flint F. The relationship of stress, competitive anxiety, mood state, and social support to athletic injury. *J Athl Train* 1996;31:296–9.
- 33 Houston JM, Harris PB, Moore R, et al. Competitiveness among Japanese, Chinese, and American undergraduate students. *Psychol Rep* 2005;97:205–12.
- 34 Jones JW, Neuman G, Altmann R, et al. Development of the sports performance inventory: A psychological measure of athletic potential. *J Bus Psychol* 2001;15:491–503.
- 35 Remor E. Propuesta de UN Cuestionario Breve para La Evaluación de la Competitividad en El Ámbito Deportivo: Competitividad-10. *Revista de Psicología Del Deporte* 2007;16:167–83.
- 36 Horvath P, Zuckerman M. Sensation seeking, risk appraisal, and risky behavior. *Personality and Individual Differences* 1993;14:41–52.
- 37 Byrnes JP, Miller DC, Schafer WD. Gender differences in risk taking: A meta-analysis. *Psychological Bulletin* 1999;125:367–83.
- 38 Morrongiello BA, Rennie H. Why do boys engage in more risk taking than girls? the role of Attributions, beliefs, and risk appraisals. *J Pediatr Psychol* 1998;23:33–43.
- 39 Hillier LM, Morrongiello BA. Age and gender differences in school-age children's appraisals of injury risk. *J Pediatr Psychol* 1998;23:229–38.
- 40 Latorre Román PÁ, Cámara Pérez JC, Pantoja Vallejo A. Effects of an education program for safety in sport in secondary school students. *Retos* 2015:5–8.
- 41 DiLillo D, Potts R, Himes S. Predictors of children's risk appraisals. *Journal of Applied Developmental Psychology* 1998;19:415–27.
- 42 Ajcardi R, Therme P. Sensation seeking, affective profile, and sports behavior in the prediction of risk. *Int J Sport Psychol* 2008;39:356–72.
- 43 Gaziantep University, School of Physical Education and Sports, Gaziantep, Turkey, Güler B, Yıldız ME, et al. Investigation of sport rock climbers' Handgrip strength. *Jbe* 2015;11:55–71.
- 44 Little H. Children's Risk-Taking behaviour: implications for early childhood policy and practice. *International Journal of Early Years Education* 2006;14:141–54.
- 45 Hansen Sandseter EB. Categorising risky play—how can we identify Risk-Taking in children's play *European Early Childhood Education Research Journal* 2007;15:237–52.
- 46 Jernbro C, Bonander C, Beckman L. The association between disability and unintentional injuries among adolescents in a general education setting: evidence from a Swedish population-based school survey. *Disabil Health J* 2020;13:100841.
- 47 Finch C. A new framework for research leading to sports injury prevention. *J Sci Med Sport* 2006;9:3–9.
- 48 González-Gálvez N, Ribeiro J, Mota J. Metabolic syndrome and cardiorespiratory fitness in children and adolescents: the role of obesity as a mediator. *J Pediatr Endocrinol Metab* 2021;34:1031–9.
- 49 Cole TJ, Donnet ML, Stanfield JP. Weight-for-height indices to assess nutritional status—a new index on a slide-rule. *Am J Clin Nutr* 1981;34:1935–43.
- 50 Léger LA, Mercier D, Gadoury C, et al. The multistage 20 metre shuttle run test for aerobic fitness. *J Sports Sci* 1986;6:93–101.
- 51 Ruiz JR, España-Romero V, Ortega FB, et al. Hand span influences optimal grip span in male and female teenagers. *The Journal of Hand Surgery* 2006;31:1367–72.
- 52 González-Gálvez N. Data from: the influence of physical fitness components on personality factors and risk perception of children and adolescents: a cross sectional study. *Dryad Digital Repository* 2022.
- 53 Cohen J. statistical power analysis for the behavioral Sciences. In: Routledge. 2013.
- 54 Baron RM, Kenny DA. The moderator–mediator variable distinction in social psychological research: conceptual, strategic, and statistical considerations. *J Pers Soc Psychol* 1986;51:1173–82.
- 55 Hernández Mendo A, Villaseñor A, Pastrana Brincones J, et al. SAGT: new software for Generalizability analysis. *Revista Iberoamericana de Psicología Del Ejercicio y El Deporte* 2016;11:77–89.
- 56 Morrongiello BA, Lasenby-Lessard J. Psychological determinants of risk taking by children: an integrative model and implications for interventions. *Inj Prev* 2007;13:20–5.

- 57 Marsh HW, Martin AJ, Jackson S. Introducing a short version of the physical self description questionnaire: new strategies, short-form evaluative criteria, and applications of factor analyses. *J Sport Exerc Psychol* 2010;32:438–82.
- 58 Balsalobre FJB, Sánchez GFL, Suárez AD. Relationships between physical fitness and physical self-concept in Spanish adolescents. *Procedia - Social and Behavioral Sciences* 2014;132:343–50.
- 59 Carroll B, Loumidis J. Children's perceived competence and enjoyment in physical education and physical activity outside school. *European Physical Education Review* 2001;7:24–43.
- 60 Hellín Gómez P, Moreno Murcia JA, Rodríguez García PL. Relación de la Competencia Motriz Percibida con La Práctica Físico-Deportiva. *Revista de Psicología Del Deporte* 2006;15:219–31.
- 61 Moreno JA, Moreno R, Cervelló E. Relación del Autoconcepto Físico con Las Conductas de Consumo de alcohol Y Tabaco en Adolescentes. *Adicciones* 2009;21:147.
- 62 Pesce C, Masci I, Marchetti R, et al. When children's perceived and actual motor competence mismatch: sport participation and gender differences. *J Mot Learn Dev* 2018;6:S440–60.
- 63 Plumert JM. Relations between children's overestimation of their physical abilities and accident proneness. *Developmental Psychology* 1995;31:866–76.
- 64 Schroth ML. A comparison of sensation seeking among different groups of athletes and Nonathletes. *Personality and Individual Differences* 1995;18:219–22.
- 65 Reio TG, Choi N. Novelty seeking in adulthood: increases accompany decline. *J Genet Psychol* 2004;165:119–33.
- 66 Hartman ML, Rawson HE. Differences in and correlates of sensation seeking in male and female athletes and Nonathletes. *Personality and Individual Differences* 1992;13:805–12.
- 67 Franken RE, Hill R, Kierstead J. Sport interest as predicted by the personality measures of competitiveness, mastery, Instrumentality, Expressivity, and sensation seeking. *Personality and Individual Differences* 1994;17:467–76.
- 68 Comín Bertrán E, Torrubia Beltri R, Mor Sancho J. Actitudes Y Consumo de alcohol, Tabaco Y Ejercicio en Escolares. *Gaceta Sanitaria* 1998;12:255–62.
- 69 Johnson HD, Sholcosky D, Gabello K, et al. Sex differences in public Restroom Handwashing behavior associated with visual behavior prompts. *Percept Mot Skills* 2003;97(3 Pt 1):805–10.
- 70 Schwebel DC, Barton BK. Contributions of multiple risk factors to child injury. *J Pediatr Psychol* 2005;30:553–61.
- 71 Moreno-Murcia JA, Belando N, Huéscar E, et al. Social support, physical exercise and life satisfaction in women. *Revista Latinoamericana de Psicología* 2017;49:194–202.
- 72 Silva MC, Malina RM. Children and youth in organized sports. 2004.
- 73 Greening L, Stoppelbein L, Chandler CC, et al. Predictors of children's and adolescents. *Journal of Pediatric Psychology* 2005;30:425–35.
- 74 Latorre Román PÁ, Cámara Pérez JC, Pantoja Vallejo A. n.d. Percepción del Riesgo en Las Actividades Físico Deportivas Escolares (risk and accident perception in sport and physical activities in school). *Retos*:93–7.
- 75 Harris MB, Miller KC. Gender and perceptions of danger. *Sex Roles* 2000;43:843–63.
- 76 Flieh SM, Miguel-Berges ML, Huybrechts I, et al. Associations between food portion sizes, insulin resistance, Vo2 Max and metabolic syndrome in European adolescents: the HELENA study. *Nutr Metab Cardiovasc Dis* 2022;32:2061–73.
- 77 Lai JCL. Dispositional optimism buffers the impact of daily Hassles on mental health in Chinese adolescents. *Personality and Individual Differences* 2009;47:247–9.
- 78 Craft LL, Pfeiffer KA, Pivarnik JM. Predictors of physical competence in adolescent girls. *Journal of Youth and Adolescence* 2003;32:431–8.
- 79 Rodríguez-Ayllon M, Cadenas-Sánchez C, Esteban-Cornejo I, et al. Physical fitness and psychological health in overweight/obese children: A cross-sectional study from the ActiveBrains project. *J Sci Med Sport* 2018;21:179–84.
- 80 Malina RM, Beunen GP, Classens AL, et al. Fatness and physical fitness of girls 7 to 17 years. *Obes Res* 1995;3:221–31.
- 81 Schnell A, Mayer J, Diehl K, et al. Giving everything for athletic success! – sports-specific risk acceptance of elite adolescent athletes. *Psychology of Sport and Exercise* 2014;15:165–72.
- 82 García Tascón M, Magaz González AM, Alías García A, et al. *La seguridad deportiva a debate*. Madrid: Dykinson, 2020.
- 83 López-Araújo B, Segovia A. Factores Explicativos de la Accidentabilidad en Jóvenes: UN Análisis de la Investigación. *Revista de Estudios de Juventud, ISSN* 2007;79:75.
- 84 Kern L, Geneau A, Laforest S, et al. Risk perception and risk-taking among Skateboarders. *Safety Science* 2014;62:370–5.
- 85 Peper JS, Braams BR, Blankenstein NE, et al. Development of Multifaceted risk taking and the relations to sex steroid hormones: A longitudinal study. *Child Dev* 2018;89:1887–907.
- 86 Latorre Román PÁ, Pantoja Vallejo A. Design and validation of a scale of perception of risk in school Físico-Deportivas activities. *Retos* 2015:25–9.
- 87 Robinson LE, Stodden DF, Barnett LM, et al. Motor competence and its effect on positive developmental Trajectories of health. *Sports Med* 2015;45:1273–84.
- 88 Sandseter EBH, Kennair LEO. Children's risky play from an evolutionary perspective: the anti-Phobic effects of thrilling experiences. *Evol Psychol* 2011;9:257–84.
- 89 Pate RR, Dowda M, Dishman RK, et al. Change in children's physical activity: predictors in the transition from elementary to middle school. *Am J Prev Med* 2019;56:e65–73.
- 90 Santos R, Mota J, Santos DA, et al. Physical fitness Percentiles for Portuguese children and adolescents aged 10–18 years. *J Sports Sci* 2014;32:1510–8.
- 91 Ortega FB, Artero EG, Ruiz JR, et al. Physical fitness levels among European adolescents: the HELENA study. *Br J Sports Med* 2011;45:20–9.
- 92 Kemper HCG, Twisk JWR, van Mechelen W. Changes in aerobic fitness in boys and girls over a period of 25 years. data from the Amsterdam growth and health longitudinal study Revisited and extended. *Pediatr Exerc Sci* 2013;25:524–35.
- 93 Janz KF, Dawson JD, Mahoney LT. Tracking physical fitness and physical activity from childhood to adolescence: the Muscatine study. *Med Sci Sports Exerc* 2000;32:1250–7.
- 94 McEwan D, Boudreau P, Curran T, et al. Personality traits of high-risk sport participants: A meta-analysis. *Journal of Research in Personality* 2019;79:83–93.
- 95 Latorre PA, Pantoja A. Design and validation of a propensity to sports accident questionnaire (PAD-22). *Cuadernos de Psicología Del Deporte* 2013;13:51–62.
- 96 Castanier C, Le Scanff C, Woodman T. Beyond sensation seeking: affect regulation as a framework for predicting risk-taking behaviors in high-risk sport. *J Sport Exerc Psychol* 2010;32:731–8.
- 97 Barrett E, Martin P. *Extreme: Why Some People Thrive at the Limits*. Oxford University Press, 2014.
- 98 Sobera A, Sobera M, Kleszyk K. Foot and ankle deformity in young acrobatic and artistic gymnasts. *Human Movement* 2015;16.
- 99 Tok S. The big five personality traits and risky sport participation. *Soc Behav Pers* 2011;39:1105–11.
- 100 Merritt CJ, Tharp JJ. Personality, self-efficacy and risk-taking in Parkour (free-running). *Psychology of Sport and Exercise* 2013;14:608–11.
- 101 Morrongiello BA, Stewart J, Pope K, et al. Exploring relations between positive mood state and school-age children's risk taking. *J Pediatr Psychol* 2015;40:406–18.
- 102 Gambetti E, Giusberti F. Anger and everyday risk-taking decisions in children and adolescents. *Personality and Individual Differences* 2016;90:342–6.
- 103 Fuster J, Elizalde B. Riesgo Y Actividades Físicas en El Medio natural: UN Enfoque multidimensional. *Apunts: Educación Física y Deportes* 1995;41:94–107.
- 104 Britton U, Issartel J, Symonds J, et al. What keeps them physically active? predicting physical activity, motor competence, health-related fitness, and perceived competence in Irish adolescents after the transition from primary to second-level school. *Int J Environ Res Public Health* 2020;17:2874.